A Guide to **Car Seat Safety**



Boston Children's Hospital

Car Seat Program
300 Longwood Avenue, Boston, MA 02115
617-355-7332
bostonchildrens.org/injury
injurypreventionprogram@childrens.harvard.edu

The Importance of Safety

Motor vehicle injuries are the number one cause of death for children from birth to 1.8 years old. Proper use of car seats and seat belts reduces serious and fatal injuries by more than 50 percent.

The Massachusetts Child Passenger Safety law passed in April of 2008 requires children to ride in a federally approved child passenger restraint that is properly fastened and secured until they are 8 years old or 4 feet 9 inches tall.

The long-term goal of the Car Seat Program at Boston Children's Hospital is that every child leaving Children's in a motor vehicle is properly restrained in a car seat. For more information, call the Car Seat Program at 617-355-7332.

Toddlers

Booster seats



- In rear-facing seats, the harness straps must be secured at or below the shoulders. The harness clip must be secured across the child's chest at ampit level.
- The harness straps must be snug: only one finger should fit under the harness strap at the collarbone.
- (2) Never place an infant seat in the front passenger seat. In the event of a car crash, an air bag could severely injure or kill an infant in a rear-facing seat
- A child may be moved into the forward-facing position when he or she is 2 years of age or has reached the maximum rear-facing limit on the car seat.
- The harness straps must be secured at or above the child's shoulders. The harness clip must be secured across the child's chest at armpit level.
- The harness straps must be snug: only one finger should fit under the harness strap at the collarbone.
- Belt-positioning booster seats are generally used for children between 4 and 8 years old, who weigh between 4 and 80 pounds. All children who have graduated from front-facing seats must be in a booster seat until they are 8 years old or 4 feet 9 inches tall, according to Massachusetts law.
- Children must use a lap and shoulder belt white riding in a booster seat.
- ⚠ Booster seats help a seat belt fit your child properly; the shoulder belt will go across the child's shoulder instead of the neck, and the lap belt will go across the hips and not the stomach.
- The American Academy of Pediatrics recommends that children 12 years of age and under sit in the rear seat of the vehicle and use an appropriate car seat or a lap and shoulder belt.

Safety belt fit test

A child is ready to use a lap and shoulder belt when he or she can:

(A) sit all the way back against the vehicle seat

- A comfortably bend their knees at the edge of the seat
- Near the shoulder belt comfortably but firmly across their shoulder, between the neck and the arm
- wear the lap belt comfortably but firmly over their hips
- (A) remain in this position for the entire trip

Special-needs car seats

Car beds are designed for infants who, for medical reasons, must travel lying down. When positioning an infant in a car bed, the infant's head must be facing the center of the vericle away from the side door.

The Hippo convertible seat from Snug Seat Inc. is designed for infarts and todders in hip casts in clicking for infarts and todders in hip casts in clicking hip pages and splittle. It is used rearraing for children from 5 to 33 pounds and forward-facing for children from 5 to 40 pounds and forward-facing from 20 to 65 pounds. The vehicle of hip cast must be included in the children with the children of the cast must be included in the

crisis swight.

For children unable to use a Hippo seat, the E-Z-ON®
Vest from E-Z-ON Products Inc. offers an alternative.
There are two types of E-Z ON Vests. The E-Z ON
Upright Vest is designed for use by children who
have behavioral concilions or poor trunk control. The
Modified E-Z-ON Vest is designed for children who
must ravel lying down.

Visit bostonchildrens.org/injury for more information. Car seat information for families living in states other than Massachusetts is available at iihs.org/laws/childrestraint.aspx.

Maintenance and installation of ALL seats

Car seats should be installed using the LATCH (lower anchors and tethers for children) or the vehicle seat belt. Vehicles manufactured after September 2002 have the LATCH installed in at least two positions in the rear.

Car seats should only be used for six years from the date of manufacture. The date of manufacture, model number and type can be found on a label on the car seat. Parents should be advised to send in their registration cards or register their seat ordine in case the car seat is recalled.

For more information on where you can get your car seat installed, call Boston Children's Car Seat Program at 617-355-7332, email at 617-355-7332, email injurypreventionprogram@childrens.harvard.edu or visit bostonchildrens.org/injury



TPA_20130125_CorSeutSaletyGuide-indd 1